

PolyMem[®]

The Ideal Dressing

Multifunctional PolyMem is a unique dressing that meets every criterion wound experts require of an ideal dressing.

Additionally, it has beneficial qualities for wound healing beyond those specifically identified.

INTRODUCTION

References to the “Ideal Dressing” date at least as far back as 1954, when John T. Scales published (in the British Journal of Industrial Medicine) a list of 10 attributes an ideal all-purpose dressing should possess.¹ The list has been modified and additional qualities have been added as researchers have learned more about how wounds heal.^{2,3,4}

Wound management courses teach that no one dressing category can meet the needs of all types of wounds from initiation of treatment to complete wound closure.^{1,4} Conventional wisdom says that because no one dressing can meet all of these criteria, clinicians must prioritize the needs of each patient, compromising on less important needs to choose the best dressing for each wound at each stage of healing.^{1,3,4}

This document examines each of the attributes that are currently considered important in an ideal dressing, as indicated by more recent literature.^{2,3,4} The conclusion is that the PolyMem family of dressings* uniquely addresses these seemingly contradictory wound needs simultaneously.

*PolyMem is often referred to in the scientific literature as “polymeric membrane dressings or PMDs”

Experts^{1,2,3,4} state an ideal dressing should:	PolyMem[®] dressings have unique attributes which make them able to meet and exceed every criterion wound experts state an ideal dressing should meet.
Maintain a moist wound environment^{2,3,4}	PolyMem dressings maintain a moist wound environment. ⁵ Because PolyMem is hydrophilic, it draws moisture from the body into the wound bed to help hydrate dry areas of the wound. ^{6,7,8,9}
Be absorbent (manage excess exudate)^{1,3,4}	PolyMem absorbs excess wound fluid and locks it into the dressings to balance moisture in the wound bed. ^{4,5,10,11,12,13,14,15,16,17,18}
Exhibit effective wound cleansing activity⁴ Remove chronic wound fluid and necrotic tissue^{3,4} Facilitate autolytic debridement^{3,4}	PolyMem provides such an effective built-in continuous wound cleansing system that even narrow deep tunnels are cleaned atraumatically. ^{6,7,10,15,19,20,21,22,23} PolyMem facilitates autolytic debridement by pulling enzyme-rich fluid from the body into the wound bed and maintaining a moist environment. ^{6,23} The dressings' surfactant loosens the bonds between the slough and wound bed. ^{4,23} The glycerin and hydrophilic membrane draw moisture into the wound bed, which floats the loosened slough. ⁵ The super-absorbent in the dressing pulls the moisture, with the slough, onto and into the dressings to be atraumatically discarded at dressing changes. ⁵
Protect the periwound from maceration⁴	PolyMem helps prevent maceration, ^{4,22,24,25} even under offloading devices and compression. ^{11,12,20}
Minimize disruption of the wound bed⁴	PolyMem continually cleanses wounds. ^{19,47} Additional cleansing is not usually necessary at dressing changes; this minimizes disruption of fragile newly-formed granulation tissue. ^{4,5,13,14,17,22,24,26,27,28,29,30,31}
Support brisk granulation and wound closure⁴	PolyMem consistently supports brisk healing. ⁴⁶ Wounds granulate and can close quickly. Recurrence is rare. ^{6,13,18,21,22,23,24,25,29,32,33,34,35,36,37,38}
Reduce or eliminate pain at the wound site³	PolyMem has the ability to subdue the nociceptor response, which can greatly reduce or even eliminate persistent wound pain. ^{4,6,13,17,18,20,22,25,26,29,30,31,35,38,39,40,41,42,43}
Protect the wound from further trauma²	PolyMem dressings are soft and flexible, providing comfortable cushioning to the wound area, and PolyMem use also usually completely eliminates traumatic wound cleansing at dressing changes. ^{6,18,28,44}
Provide thermal insulation^{2,3,4}	Wound cooling can slow healing. ^{3,36} Because rinsing at dressing changes is rarely needed, PolyMem use diminishes wound cooling. ²⁸ The PolyMem membrane insulates to keep wounds warm. ^{10,25,36}

Control wound odor⁴	PolyMem dressings' formulation provides natural odor control. ^{5,18}
Be permeable to gas and water vapor¹	The PolyMem membrane and the semipermeable outer film permit controlled exchange of oxygen and water vapor. ^{9,15}
Act as a bacterial barrier^{1,2,3,4}	All PolyMem secondary dressings have a semipermeable outer film which protects against microbial contamination. ^{9,25,28,38}
Provide antimicrobial protection^{1,4}	PolyMem Silver [®] dressings harness the antimicrobial properties of silver to prevent the dressings from harboring microbes. ^{4,10,15,16,21,22,25,28}
Be non-toxic and non-allergenic^{1,2,4}	<p>Independent researchers tested the relative toxicity of wound cleansers and found that PolyMem's surfactant was the least cytotoxic; it was less toxic than saline.³⁵</p> <p>Other silver dressings leach cytotoxic silver into the wound bed, but the silver in PolyMem Silver acts from within the dressing, allowing new tissue and fibroblasts to proliferate.^{16,25,28}</p> <p>PolyMem is non-toxic, is not made with latex, and does not irritate patients' skin.^{9,11,15,21,27,35,45,46}</p>
Be comfortable, even when used to fill tunneling, undermining, or dead space^{2,3}	In addition to being comfortable to the wearer, ⁴⁷ PolyMem WIC [®] and PolyMem WIC Silver Rope can reduce persistent wound pain when used in tunneling, undermining and dead space. ^{4,12,22,25,26,28,31}
Be conformable, even over joints (fingers can flex while dressed)^{1,2,4}	PolyMem is stretchy and conformable, ^{42,48} making it an ideal dressing choice for use over joints. ^{7,36} The PolyMem family of dressings includes specially designed finger and toe dressings for added convenience. ^{20,27,31}
Not allow tissue to grow into dressing¹	The PolyMem membrane is specially designed to prevent tissue from growing into the dressing; this is a problem with so many other dressings. ^{15,34}
Not cause pain or tissue damage on dressing removal^{2,3,4}	Because the dressings are nonadherent and tissue does not grow into PolyMem, dressing changes are comfortable and atraumatic. ^{4,6,13,15,18,22,26,30,34,42,44,45}

<p>Not leave particles or toxic contaminants in the wound bed^{2,4}</p>	<p>PolyMem dressings are engineered to remain intact, even when fully saturated.^{21,28,42,45} PolyMem WIC Silver Rope is reinforced with surgical mesh to prevent breakage, even when inserted into deep, narrow tunnels. ^{13,21,22}</p> <p>The silver in PolyMem Silver kills microbes while remaining locked in the dressing, thus minimizing leaching of cytotoxic silver into the wound bed.^{16,21,22,32}</p>
<p>Stay in place⁴</p>	<p>PolyMem dressings are proven to stay in place, even when patients do not adhere to the treatment plan.^{6,7,18,25,36,44,47,48}</p>
<p>Come in various shapes and sizes³</p>	<p>The PolyMem family of dressings includes a wide variety of shapes and sizes to meet every wound need.^{9,10,13,23,47}</p>
<p>Require only infrequent dressing changes²</p>	<p>PolyMem dressings are changed only when “indicated” by the level of exudate drawn into the dressing.⁴⁵ While dressing changes may be as frequent as twice daily initially, continuous wound cleansing and moisture balancing allow PolyMem to safely remain in place up to one week after exudate levels stabilize.^{13,20,23,32,33}</p>
<p>Be nonadherent to blood clots and granulating surfaces¹</p>	<p>The wound contact surface of a PolyMem dressing becomes a soft, conformable, nonadherent gel when activated by wound fluid.^{6,11,22,42} In addition, PolyMem continuously draws fluid into the wound bed, which keeps the dressing from drying onto the wound surface.^{6,18,27,28,31}</p>
<p>Be sterilizable¹</p>	<p>PolyMem dressings are gamma irradiated. Packages are guaranteed sterile until opened or expired.</p>
<p>Have a long shelf life²</p>	<p>PolyMem dressings stored at room temperature are guaranteed for up to 5 years from manufacturing date.</p>
<p>Be quick and easy to use⁴</p>	<p>PolyMem is so quick and easy to use^{20,21,32,47} that when guided by health care professionals, patients and families can often perform the dressing changes themselves.^{5,6,12,14,15,18,22,23,26,27,33,35,44}</p>
<p>Be cost effective^{1,2}</p>	<p>The modest initial dressing price, decreased time and supplies needed for dressing changes, and quick healing work together to make its use extremely cost effective.^{21,32,47,49,50} Facilities often make PolyMem their dressing of choice.^{6,11,20,23,33,35,38,47}</p>

<p>If self-adhesive, forms an effective seal while being removable without skin stripping⁴</p>	<p>PolyMem Shapes are the preferred dressings for skin tears in many Post Acute Care facilities due to their excellent water resistant adhesion and atraumatic removal.^{47,51}</p>
<p>Decrease bleeding⁴</p>	<p>In nasal surgery, PolyMem dressings provided excellent perioperative bleeding control and performed better than other dressings.⁵² PolyMem was found to make a safe and effective intranasal dressing for patients undergoing control of epistaxis.⁵³</p> <p>The PolyMem membrane structure provides high surface area and fast absorption of liquids; both of these properties help facilitate hemostasis. In addition, the PolyMem membrane is thought to selectively absorb the watery portion of exudate, thus concentrating clotting factors.</p>

Multifunctional PolyMem is a unique dressing which meets all of these criteria, and it has attributes *beyond* the ones wound experts identified which help heal wounds.

PolyMem also helps to:

- Improve the nutrition of ischemic wounds^{6,54,55}
- Focus the inflammation, which is an integral part of the chronic wound state, into the damaged tissues while reducing the spread into the surrounding tissues through nociceptor inhibition.^{15,20,26,38,40,41,56,57}
- Decrease edema, bruising and pain associated with tissue injury^{6,9,26,33,39,46,57}
- Strengthen the scar (due to continued nociceptor inhibition) during the remodeling phase of wound healing.^{18,37,58}
- Indicate when dressings require changing (a color change is visible through the semipermeable outer layer of the dressing), minimizing disruption of the wound bed.⁴⁴

Most wound courses and textbooks teach that the dressing choice must be altered at various stages of healing, depending upon the current condition of the wound. However, PolyMem provides all of the benefits of an ideal dressing.^{1,3,4} PolyMem can even add moisture to dry areas of a wound while removing excess moisture from highly exudative

areas, making it suitable for use over exposed deep structures like tendons and bones.²⁸ PolyMem can be used at any stage of healing and through all stages of healing for virtually all wounds.

PolyMem in either the standard or the silver configurations can be used to continuously cleanse infected wounds.⁹ The cause of the infection should be addressed appropriately.⁹ PolyMem Silver dressings, which contain small particle silver, have been proven to kill over 99% of the populations of representative organisms.³² All PolyMem dressings pull exudate, drainage, dirt, slough, and fibrin into the dressing matrix; PolyMem Silver dressings also help manage bioburden and kill microbes in contact with the dressing.^{24,32} Rather than placing silver directly into the wound bed, the silver in PolyMem Silver is built into the dressing to help eliminate the bioburden and microbes in the dressing, doing potentially less damage to the healing wound.^{16,21,28,37}

Wound management must be affordable, clinically effective, and convenient to both the patient and care providers. PolyMem use promotes patient independence and increased quality of life.^{12,13,20,21,23,36,38} Implementing protocols which use PolyMem dressings alone has proven to be more clinically effective, cost effective, and convenient than implementing conventional wound protocols.^{10,23,31,36,49-51,59-61}

References

1. Scales JT. Some Problems of Wound Dressings. *Occup Med (Lond)*. 1954 Jul 1;4(2):69-72.
2. Jones V, Grey J, Harding K. Wound dressings. *BMJ*. 2006 Apr 1;332(7544):777-80.
3. Baranoski S, Ayello EA. *Wound care essentials: practice principles*. 2nd ed. Philadelphia: Lippincott Williams & Wilkins; 2008.
4. Thomas S. *Surgical Dressings and Wound Management*. 2nd ed. Hinesburg VT: Kestrel Health Information; 2012. 708 p.
5. Agathangelou C. Deep ulcer on charcot foot closed after treatment with polymeric membrane silver cavity dressing*. Poster presented at the 3rd congress of the World Union of Wound Healing Societies June 4-8, 2008. Toronto, Ontario Canada.
6. Agathangelou C. Painless Autolytic Debridement on 250 Chronic Wounds by Using Polymeric Membrane Dressings. 4th Congress of the World Union of Wound Healing Societies Po #241 (WUWHS) September 2-6, 2012 Pacifico Yokohama, Yokohama, Japan
7. Benskin L. Chronic Wounds Close Using Various Configurations of Polymeric Membrane Dressings Despite Fear-Generated Non-Compliance. 3rd Congress of the World Union of Wound Healing Societies Poster #PF422 June 4-8, 2008 Toronto, Ontario Canada.
8. Benskin L. Deep Ulceration Treated with Polymeric Membrane Dressings Until Complete Wound Closure. 3rd Congress of the World Union of Wound Healing Societies Poster #PF422 June 4-8, 2008 Toronto, Ontario Canada
9. Hess CT. *Clinical Guide to Skin and Wound Care*. 7th edition. Lippincott Williams & Wilkins; 2012.

10. Seemann P. Stalled infected diabetic foot wounds respond rapidly to various polymeric membrane dressing* types. Poster presented at the 22nd annual (SAWC) Symposium on Advanced Wound Care & WHS meeting, April 26-29, 2009. Dallas, TX.
11. Curtin, C. The new diabetic foot ulcer standard of care: Instant total contact cast plus polymeric membrane dressings. Poster presented at 9th annual Desert Foot: High Risk Diabetic Foot Conference. Nov 14-16, 2012. Phoenix, AZ.
12. Hjalmarsson M. Diabetic foot saved from amputation with Polymeric membrane dressings. 19th Conference of the European Wound Management Association, Poster #228. May 20-22, 2009, Helsinki, Finland.
13. Yip WCA, Wong M, Yu OHI, Choi YN, Tang SF, Hui HK, May MK. Extensive, Amputation threatening, diabetic foot ulcer closed rapidly with multifunctional silver polymeric membrane dressings. Australian Wound Management Association, Poster #59 March 22-24, 2010, Perth, Western Australia
14. Benskin L. Combining Email Wound Consulting and Polymeric Membrane Dressings to Dramatically Improve Quality of Life. 22nd Annual (SAWC) Symposium on Advanced Wound Care & WHS Meeting Poster #R003 April 26-29, 2009 Dallas, TX USA.
15. Smith D, Houle H, Searls L. Hypergranulation Tissue, A Challenging Problem with an Easy Solution. WOW Wild On Wounds Conference Poster #21 September 11-14, 2012 Las Vegas, Nevada.
16. Burd A, Kwok CH, Hung SC, Chan HS, Gu H, Lam WK, Huang L. A comparative study of the cytotoxicity of silver-based dressings in monolayer cell, tissue explant, and animal models. Wound Repair Regen 2007;15:94-104.
17. Langemo DK, Black J; National Pressure Ulcer Advisory Panel. Pressure ulcers in individuals receiving palliative care: a National Pressure Ulcer Advisory Panel white paper. Adv Skin Wound Care 2010;23:59-72.
18. Denyer J. Polymeric membrane dressings in the management of infants with epidermolysis bullosa (EB). 19th Conference of the European Wound Management Association Poster #100 May 20-22, 2009, Helsinki, Finland.
19. Benskin LLL. PolyMem Wic Silver Rope: A Multifunctional Dressing for Decreasing Pain, Swelling, and Inflammation. Advances in Wound Care. 2012 Feb;1(1):44-7.
20. Selekof JF, Shutt D, Roney A. Diabetic foot ulcers: multicenter case studies using silver polymeric membrane dressings to promote wound healing. Symposium on Advanced Wound Care Fall, Poster #CS-19 September 16 -19, 2009, Washington, D.C. USA
21. Nair HKR, Har MY. Utilisation of Multifunctional Dressing on Stagnate, Draining Neuropathic Wound Using Polymeric Membrane Silver Dressing. Australian Wound Management Association conference, 18-21 March 2012 Sydney, Australia.
22. Tamir J. Mesh reinforced silver rope dressing for acute infected cavity wounds. Symposium on Advanced Wound Care (SAWC) SPRING Poster #CS-118 April 17-20, 2010 Orlando, FL USA
23. Reaves K. Silver polymeric membrane dressings*: An economical choice for resolving diabetic wounds. Poster presented at the 21st annual (SAWC) Symposium on Advanced Wound Care & WHS meeting, April 24-27, 2008. San Diego, CA.
24. Benskin L. Diabetic foot salvaged, wounds closed in only two months using polymeric membrane cavity filler* and polymeric membrane dressings*. Poster presented at 39th annual meeting of the Wound Ostomy Continence Nurses Society (WOCN), June 10-13, 2007. Salt Lake City, UT.
25. Agathangelou C. Huge sacral pressure ulcer closed in four months using silver polymeric membrane cavity filler and dressings. NPUAP 11th Biennial Conference, Poster #40 February 27-29, 2009, Arlington, VA USA
26. Agathangelou C. Patients with Chronic Wound Pain Stop Hurting, Why? European Wound Management Association (EWMA) Poster #348 May 25-27, 2011 Brussels, Belgium
27. Benskin L. Excellent healing of pediatric wounds using polymeric membrane dressings. 41st Annual WOCN Conference, Poster #3228. June 6 -10, 2009, St. Louis, MO USA

28. Benskin L. Extensive tunneling lower leg wounds with exposed tendons closed quickly using various polymeric membrane dressing configurations. 23rd Annual Clinical Symposium on Advances in Skin & Wound Care, Poster #41 October 26-30,2008, Las Vegas, NV USA
29. Lee BY. The diabetic foot: a comprehensive approach. In: Lee BY, editor. The wound management manual. New York: McGraw-Hill; 2005:360-1.
30. Davies SL, White RJ. Defining a holistic pain-relieving approach to wound care via a drug free polymeric membrane dressing. J Wound Care. 2011 May;20(5):250, 252, 254 passim.
31. Vanwalleghem, G. Rapid closure of infected diabetic foot through the use of polymeric membrane cavity filler*. Poster presented at the 19th conference of the European wound management association, May 20-22, 2009. Helsinki, Finland.
32. Wilson, D. Stalled diabetic ulcer closed in six weeks using PolyMem Silver dressings. Poster presented at 39th annual meeting of the Wound Ostomy Continence Nurses Society (WOCN), June 10-13, 2007. Salt Lake City, UT.
33. Ogden J. A "New Approach" to Healing Chronic Wounds Accompanied by Dramatically Lower Cost of Wound Management. WOW Wild On Wounds Conference Poster #1 September 7-10, 2011 Las Vegas, Nevada USA
34. Schwindling R. Schon wieder eine neue wundauglage? V.A.C. therapy und Polymem - ein ester erfahrungsbericht. KCI Times 2013
35. Wilson JR, Mills JG, Prather ID, Dimitrijevic SD. A toxicity index of skin and wound cleansers used on in vitro fibroblasts and keratinocytes. Adv Skin Wound Care. 2005 Sep;18(7):373-8.
36. Agathangelou C. Injured finger salvaged using PolyMem Silver dressings to keep the wound bed clean and warm. 3rd Congress of the World Union of Wound Healing Societies, Poster #PF430 June 4-8, 2008, Toronto, Ontario Canada
37. Yastrub DJ. Relationship between type of treatment and degree of wound healing among institutionalized geriatric patients with stage II pressure ulcers. Care Manag J 2004;5:213-218.
38. Rahman S, Shokri AA. Total Knee Arthroplasty (TKA) Infections Eliminated and Rehabilitation Improved Using Polymeric Membrane Dressing Circumferential Wrap Technique: 120 Patients at 12-month Follow-up. European Wound Management Association (EWMA) Poster #398 May 15-17, 2013 Copenhagen, Denmark
39. Man D, Aleksinko P. Intra-Operative Use Followed with Post-Op Application of Polymeric Membrane Dressings Reduces Post-Op Pain, Edema and Bruising After Full Face Lift Surgery. Symposium on Advanced Wound Care (SAWC) Spring Poster #CS 071 April 17-20, 2010 Orlando, FL USA
40. Hayden JK, Cole BJ. The effectiveness of a pain wrap compared to a standard dressing on the reduction of postoperative morbidity following routine knee arthroscopy: a prospective randomized single-blind study. Orthopedics 2003;26:59-63.
41. Beitz AJ, Newman A, Kahn AR, Ruggles T, Eikmeier L. A polymeric membrane dressing with antinociceptive properties: Analysis with a rodent model of stab wound secondary hyperalgesia. The journal of pain.2004;5(1):38-47.
42. Man D, Aleksinko P. Use of Polymeric Membrane Dressings Immediately After Fractionated Facial Laser Resurfacing Procedures Improves Outcomes. Symposium on Advanced Wound Care (SAWC) Spring Poster #CS-069 April 17-20, 2010 Orlando, FL USA
43. Agathangelou C. Three years experience of treating 46 painful diabetic foot ulcers with polymeric membrane dressings. European Wound Management Association (EWMA) Poster #47 May 23-25, 2012. Vienna, Austria
44. Agathangelou C. Closing stalled heel pressure ulcers: two problem patients, one easy solution. NPUAP 11th Biennial Conference, Poster #33 February 27-29, 2009, Arlington, VA USA
45. Fowler E, Papen JC. Clinical evaluation of a polymeric membrane dressing in the treatment of dermal ulcers. Ostomy Wound Manage. 1991 Aug;35:35-8, 40-4.
46. Rodeheaver GT, Kurtz L, Kircher BJ, Edlich RF. Pluronic F-68: a promising new skin wound cleanser. Ann Emerg Med. 1980 Nov;9(11):572-6.

47. Wilson D. Skin tear healing improved through the use of polymeric membrane dressings. 3rd Congress of the World Union of Wound Healing Societies, Poster #PW725 June 4-8, 2008, Toronto, Ontario Canada
48. Benskin L. Crush injury treated with extra-thick polymeric membrane dressings until complete wound closure. 3rd Congress of the World Union of Wound Healing Societies, Poster #PW368 June 4-8, 2008, Toronto, Ontario Canada
49. Hoglin G, Melin A. Improving clinical outcome while reducing cost of treatment. Poster #108. European Wound Management Association (EWMA). May 23-15, 2012 Vienna, Austria
50. Pacheco MS. Best clinical practices, based on single dressing type, improves wound healing while reducing health care costs. Wild on Wound Conference (WOW) Poster #9 September 11-14, 2013, Las Vegas, Nevada
51. Bolhuis J, Benskin L. Evidence-based skin tear protocol yields phenomenal results. Poster #2403 40th Annual WOCN Conference. June 21-28, 2008 Orlando, FL
52. Phillips PP. Postoperative endonasal dressing with polymeric membrane material. Poster presented at: American Rhinologic Society, September 2003 - Orlando, FL
53. Phillips PP. Polymeric membrane dressing role in epistaxis control. Poster presented at: American Academy of Otolaryngology - Head and Neck Surgery Foundation, Inc. September 2003 - Orlando, FL USA
54. Yastrub, Heel Ulcer in Hospice Patient Closed Quickly Using Polymeric Membrane Dressings. Poster presented at 40th annual meeting of the Wound Ostomy Continence Nurses Society (WOCN), Orlando, FL. June 21 -25, 2008.
55. Agathangelou C. Unique dressing provides nutrients for wound closure in a profoundly malnourished patient. NPUAP 11th Biennial Conference, Poster #36 February 27-29, 2019, Arlington, VA USA
56. Wilson D. Application of Polymeric Membrane Dressings to Stage I Pressure Ulcers Speeds Resolution, Reduces Ulcer Site Discomfort and Reduces Staff Time Devoted to Management of Ulcers. Symposium on Advanced Wound Care (SAWC) SPRING Poster # CS-128 April 17-20, 2010 Orlando, FL USA
57. Benskin L. PolyMem® WIC® Silver® Rope; A Multifunctional Dressing for Decreasing Pain, Swelling, and Inflammation. Adv Wound Care (New Rochelle). 2012 Feb;1(1):44-7.
58. Harrison J. Polymeric membrane dressings bring brisk healing to bilateral pressure ulcers. NPUAP 11th Biennial Conference, Poster #32 February 27-29, 2009, Arlington, VA USA
59. Winblad R, Harvey R. A new wound care approach yields significant clinical, economic, logistical and training satisfaction. Poster #P121 European Wound Management Association (EWMA) meeting. May 26-28, Geneva Switzerland
60. Stenius M. A holistic approach together with polymeric membrane dressings closed Grade IV pressure ulcer in 2.5 months. Poster #P247 European Wound Management Association (EWMA). May 14-16, 2008 Lisbon, Portugal.
61. Pacheco MS. Best clinical practices, based on single dressing type, improves wound healing while reducing health care costs. Wild on Wound Conference (WOW) Poster #9 September 11-14, 2013, Las Vegas, Nevada


With PolyMem dressings, you no longer need to rely on multiple wound care products. PolyMem has everything you need in one easy-to-use dressing. PolyMem is indicated for virtually all wounds at every stage of healing.

PolyMem dressing are available in a wide variety of configurations, with or without silver, to meet every wound need:

- Non-adhesive pads and rolls
- Rectangular thin film or cloth adhesive bordered dressings
- Surgical bordered dressings
- Oval Shapes® dressings bordered with silicone or acrylic adhesive
- Sacral Shapes bordered dressings
- Shapes dressings for tube sites
- Extra-thick MAX® pads with or without adhesive border
- WIC® cavity filler in a variety of sizes, plus WIC Silver Rope
- Finger/Toe dressings in five sizes

SportsWrap® dressings are nonsterile products especially designed for closed injuries.

Learn more about PolyMem at www.polymem.com



PolyMem dressings are unique dressings that meet and exceed the criteria wound experts require in an ideal dressing.

PolyMem is indicated for virtually every wound clinicians may encounter.



5133 Northeast Parkway, Fort Worth, TX 76106-1822 U.S.A.
Toll Free U.S.A.: 800-POLYMEM (765-9636)
International: +1 817-900-1301
www.polymem.com

This document is meant for general information. See individual product instructions for use.

Trademarks are owned by or licensed to Ferris. The marks may be registered or pending in several countries.

© 2015-2019 Ferris Mfg. Corp.

MKL-662 R3 1119